AD-A065 088

ARMY MEDICAL RESEARCH INST OF INFECTIOUS DISEASES FR-ETC F/G 6/5 LEGIONNAIRES DISEASE BACTERIA: REPLICATION IN ALVEOLAR MACROPH-ETC(U) JAN 79 R A KISHIMOTO, M D KASTELLO, J D WHITE

UNCLASSIFIED

ADA 065088

###



















END DATE FILMED

NL

4 -79

Legionnaires' Disease Bacteria: Replication in
Alveolar Macrophages of Cynomolgus Monkeys.

R. A. KISHIMOTO, M. D. KASTELLO, J. D. WHITE, F. G. SHIREY

V. G. McGANN E. W. LARSON, AND K. W. HEDLUND

16) 3M762776A841 (1) MAR 1 1979

B

Vinited States Army Medical Research Institute of Infectious Diseases

Fort Detrick, Frederick, Maryland 21701

The views of the authors do not purport to reflect the positions of the Department of the Army or the Department of Defense.

In conducting the research described in this report, the investigators adhered to the "Guide for the Care and Use of Laboratory Animals," as promulgated by the Committee on the Revision of the Guide for Laboratory Animal Facilities and Care of the Institute of Laboratory Animal Resources, National Research Council. The facilities are fully accredited by the American Association for Accreditation of Laboratory Animal Care.

26 Jan 1979 (2)

Cleared for public release; distribution unlimited.

405 039

43

DE FILE COPY

Abstract. The interaction between normal cynomolgus monkey

alveolar macrophages and Legionnaires' disease bacteria (LDB) was

studied by transmission electron microscopy. Following ingestion of

LDB, the organisms replicated within macrophages and destroyed the

phagocytic cell.

ACCESSION	fer
NTIS DDC UNANNOTED JUSTIFICATI	
BY DISTRIBUNG	N/AVAILABILITY CODES
Dist. An	il. and/or SPECIAL
A	

A severe outbreak of pneumonia involving 182 individuals with 29 deaths occurred at an American Legion convention in Pennsylvania in the summer of 1976 (1, 2). The consistent pathologic feature noted was acute pneumonia with an infiltrate of polymorphonuclear neutrophils and macrophages (3-5). Microscopically, dense clusters of organisms which often obscured nuclear detail were observed in lung macrophages (3). Disruption of phagocytic cells accompanied by many extracellular bacteria has also been noted (5). These observations have led to the hypothesis that the organisms resist digestion and replicate within phagocytes (3, 5).

Preliminary studies in our laboratories indicate that cynomolgus monkeys (Macaca fascicularis) show clinical signs of illness following an aerosol-induced challenge of the Washington strain of Legionnaires' disease bacteria (LDB) (unpublished data). The objective of the current study was to examine by transmission electron microscopy (TEM) in vitro interactions between LDB and alveolar macrophages obtained from normal cynomolgus monkeys.

The Washington strain of LDB which is virulent for guinea pigs was cultured on Mueller-Hinton agar supplemented with 2 percent Isovitalex (6) and 1 percent hemoglobin and incubated at 37°C for 48 hours in a humid atmosphere of air containing 5 percent CO₂. Organisms were scraped off the plates, washed once in Hank's balanced salt solution (HBSS), and resuspended in sufficient Earle 199 medium supplemented with 10 percent normal cynomolgus serum so that the final suspension contained approximately 10⁸ colony forming units/ml.

Cynomolgus monkeys of both sexes, weighing 2.0-3.5 kg were used in this study. Alveolar macrophages were recovered from anesthetized,

normal monkeys by lung lavage (7). Approximately 25 x 10⁶ macrophages in 25 ml Earle 199 medium supplemented with 10 percent normal cynomolgus monkey serum were dispensed into 60 x 15 mm petri dishes (8), incubated at 37°C for 3 hours in a humid atmosphere containing 5 percent CO₂, then washed twice with HBSS to remove nonadherent cells. AntibiotTcs were excluded in this study. Macrophage cultures were inoculated with LDB at a ratio of 100 organisms per macrophage. After incubation at 37°C for 3 hours, macrophage cultures were washed three times with 25 ml of HBSS. Some cultures were processed for examination by TEM (9), while others were re-incubated in fresh medium for an additional 21 hours at 37°C. All TEM preparations were viewed with a Hitachi HU-12 electron microscope operated at 75 kV.

After the 3-hour interaction period, approximately 5 percent of the macrophages showed evidence of intracellular LDB. These cells contained 1-3 organisms which appeared to be contained within a membrane-bound cytoplasmic vesicle (Fig. 1A). Twenty-four hours later, many macrophages contained distended vacuoles filled with LDB (Fig. 1B). Multiplication of organisms appeared to be so rapid and extensive that the entire cytoplasmic compartment of some cells became filled with vesicles containing LDB (Fig. 2) and ultimately the phagocytes were destroyed. Figure 3 shows that LDB are very pleomorphic. The cell structure of the organisms resembled that of other gram-negative bacteria (Fig 3), as well as those described by Katz and Nash (10). In Fig. 3 two organisms undergoing binary fission can be seen.

Often the outcome of the interaction between invading organisms and alveolar macrophages determines the extent of infection (11). If organisms are killed by the macrophage, disease is averted; replication of the invading organism within the phagoctye causes the host to be compromised.

The lung is recognized as one of the most frequently involved organs in Legionnaires' disease (1, 3, 4), but the pathogenesis has not been elucidated. Our data indicate that LDB are not avidly phagocytized by the alveolar macrophage. However, when ingested, the organisms replicated rapidly and destroyed the phagocyte. Alveolar macrophages from normal cynomolgus monkeys would not appear to provide a defense against initial infection with this organism. This observation suggests that LDB can abrogate one of the primary defense mechanisms of the host. Additional studies are necessary to elucidate the role of other defense mechanisms such as polymorphonuclear leukocytes, specific antibody, and cell-mediated immunity.

References and Notes


- D. W. Fraser, T. R. Tsai, W. Orenstein, W. E. Parkin, H. J.
 Beecham, R. G. Sharrar, J. Harris, G. F. Mallison, S. M. Martin,
 J. E. McDade, C. C. Shepard, P. S. Brachman, the Field Investigation
 Team, N. Engl. J. Med. 297, 1189 (1977).
- Center for Disease Control, <u>Morbidity and Mortality Weekly</u>
 Report 25, 244 (1976).
- 3. J. A. Blackmon, M. D. Hicklin, F. W. Chandler, the Special Expert Pathology Panel, Arch. Pathol. Lab. Med. 102, 337 (1978).
- 4. F. W. Chandler, M. D. Hicklin, J. A. Blackmon, N. Engl. J. Med. 297, 1218 (1977).
- W. C. Winn, Jr., F. L. Glavin, D. P. Perl, J. L. Keller, T. L. Andres, T. M. Brown, C. M. Coffin, J. E. Sensecqua, L. N. Roman, J. E. Craighead, Arch. Pathol. Lab. Med. 102, 344 (1978).
- 6. Purchased from Difco Laboratories, Detroit, Mich.
- 7. M. D. Kastello, A. D. Emmert, R. F. Denson, R. A. Kishimoto,

 Am. J. Vet. Res., in press (1979).
- 8. Purchased from Falcon Plastics, Oxnard, Calif.
- R. A. Kishimoto, B. J. Veltri, P. G. Canonico, F. G. Shirey,
 J. S. Walker, <u>Infect</u>. <u>Immun</u>. 14, 1087 (1976).
- 10. S. M. Katz and P. Nash, Am. J. Pathol. 90, 701 (1978).
- E. Goldstein, W. Lippert, D. Warshauer, J. Clin. Invest. 54, 519 (1974).
- 12. We thank Robert W. Stockman and Stephen F. Little for their excellent technical support.

Figure Legends

- Fig. 1. Alveolar macrophages containing Legionnaires' disease bacteria. A. Single organism in phagocytic vesicle at 3 hours (X8000). Bar, 1.0 μ m. B. Large numbers of bacteria within a distended vesicle at 24 hours (X8,000). Bar, 1.0 μ m.
- Fig. 2. Alveolar macrophage at 24 hours. The cytoplasm is filled with numerous small vesicles containing pleomorphic forms of Legionnaires' disease bacteria (X8,000). Bar, 1.0 μm .
- Fig. 3. Cytoplasm of alveolar macrophage containing Legionnaires' disease bacteria. Note evidence of binary fission (♠), cell wall of bacterium (♠), and vesicular membrane (♠) (X30,000).

 Bar, 1.0 μm.



REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 2.	GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (end Subtitle) LEGIONNAIRES' DISEASE BACTERIA: REPLICATION IN ALVEOLAR MACROPHAGES OF CYNOMOLGUS MONKEYS		5. TYPE OF REPORT & PERIOD COVERED Interim 6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(*) R.A. KISHIMOTO, M.D. KASTELLO, J.D. WHITE, F.G. SHIREY, V.G. MCGANN, E.W. LARSON, AND		8. CONTRACT OR GRANT NUMBER(*)
W. HEDLJIND 9. PERFORMING ORGANIZATION NAME AND ADDRESS U.S. Army Medical Research Institute of Infectious Diseases SGRD-UIA-E Fort Detrick, Frederick, Maryland 21701		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 3M762776 A841-00-007
U.S. Army Medical Research and Development Command, Office of the Surgeon General Department of the Army, Washington, DC 20310 14. MONITORING AGENCY NAME & ADDRESS(II different from Controlling Office)		12. REPORT DATE 26 January 1979 13. NUMBER OF PAGES 8 15. SECURITY CLASS. (of this report) UNCLASSIFIED
		154. DECLASSIFICATION/DOWNGRADING SCHEDULE

16. DISTRIBUTION STATEMENT (of this Report)

Cleared for public release; distribution unlimited.

17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)

18. SUPPLEMENTARY NOTES

Reprints bearing assigned AD number will be forwarded upon receipt. To be submitted for publication in Infection and Immumity.

19. KEY WORDS (Continue on reverse side if necessary and identify by block number)

Legionnaires' Disease Bacteria Alveolar Macrophage Cynomolgus Monkey Transmission Electron Microscopy

20. ABSTRACT (Continue on reverse side If necessary and identify by block number)

The interaction between normal cynomolgus monkey alveolar macrophages and Legionnaires' disease bacteria (LDB) was studied by transmission electron microscopy. Following ingestion of LDB, the organisms replicated within macrophages and destroyed the phagocytic cell.

DD 1 JAN 73 1473 EDITION OF 1 NOV 65 IS OBSOLETE